

Urologist



Dr Jo Schoeman
Specialist Urologist

PATIENT INFORMATION BROCHURE

TRANS URETHRAL DE-ROOFING OF PROSTATE ABSCESS

See this live on:
vidscrip.com/urojo

Patient well-being is my first priority!

Jo Schoeman
FRACS, FCS (Urol) SA, MBChB

The Wesley Hospital
Suite 10 Level 9
Evan Thompson Building
24 Chasely street
AUCHENFLOWER QLD 4066

Ph: 07) 3371-7288
Fax: 07) 3870-5350
E-mail: jo@urojo.com.au
Emerg: 0403 044 072


www.brisbane-urologist.com.au

Trans Urethral De-roofing of Prostate Abscess

Non invasive opening of an abscess found in the prostate, Very similar to a TURP with the exception that the bladder neck is preserved

Why is it done?

- This procedure is performed when you have been diagnosed with a prostatic abscess, usually after an MRI investigation
- You would have had a history of swinging low to high grade fevers, a history of catheterization or instrumentation of the urethra, possibly a prostate biopsy.
- Not improvement on Antibiotics.
- You may possibly be very sick with High Dependency Unit admission on intravenous antibiotics
- This procedure is done where the abscess or fluid collection in the parenchyma of the prostate is in the transitional zone of the prostate and easily accessible via endoscopic technique
- It is done under a General Anaesthetic with prophylactic antibiotics in place, or appropriate Antibiotics as per your serum cultures



How is it done?

- Patients will receive a general Anaesthesia unless otherwise indicated.
- A cystoscopy is performed by placing a camera in the urethra with the help of lignocaine gel.
- The inside of the prostatic urethra and bladder is viewed for pathology and especially signs of the abscess. If any suspicious lesions are seen, a biopsy will be taken.
- The area of the prostatic urethra above the abscess is resected until the abscess is opened and drained.
- The bladder neck and urethral sphincter is preserved.
- Laser can also be utilized and is probably preferred due to lack of bleeding
- Prophylactic antibiotics will be given to prevent any infections.

What can go wrong?

- Any anaesthesia has its risks and the anaesthetist will explain this to you..
- You will wake up with a catheter in your urethra and bladder. This will remain in the bladder for 1-3 days depending on the technique used and incidence of post-operative bleeding. And until signs of sepsis have cleared
- You may have a continuous bladder irrigant running in and out of your bladder to prevent clot formation.
- Lower abdominal discomfort for a few days
- NB! Each person is unique and for this reason symptoms vary!

What next?

- You will be hospitalised until all signs of sepsis have cleared
- You will have a trial without catheter as soon as your urine is clear.
- You will be discharged as soon as you can completely empty your bladder.
- You may initially suffer from urge incontinence and will improve within the next 6 weeks.
- Allow for 6 weeks for stabilization of symptoms.
- You may have a change in ejaculate volume and could even suffer retrograde ejaculation.
- There may be some blood in your urine. You can remedy this by drinking plenty of fluids until it clears.
- A ward prescription will be issued on your discharge, for your own collection at any pharmacy
- A follow-up appointment will be scheduled for 6 weeks. Should your pathology be worrisome, you will be contacted for an earlier appointment.
- Don't hesitate to ask me if you have any queries
- **DON'T SUFFER IN SILENCE, OR YOU WILL SUFFER ALONE!**

