

Complications:

- Any anaesthesia has its risks and the anaesthetist will explain this to you.
- Complications may vary depending on the initial surgery done
- Urine leak
- Ureteric stent X 6 weeks
- Infection
- Stent irritation
- Strictureing at the site of surgery requiring dilatations, permanent stent or re-do surgery
- NB! Each person is unique and for this reason symptoms vary!

Narrowing of passage
due to strictures



Normal



Moderate



Severe



Total

Jo Schoeman
FRACS, FCS (Urol) SA, MBChB

The Wesley Hospital
Suite 10 Level 9
Evan Thompson Building
24 Chasely street
AUCHENFLOWER QLD 4066

Ph: 07) 3371-7288
Fax: 07) 3870-5350
E-mail: jo@urojo.com.au
Emerg: 0403 044 072

www.brisbane-urologist.com.au

Urologist



Dr Jo Schoeman
Specialist Urologist

**PATIENT
INFORMATION
BROCHURE**

***URETERIC
INJURY***

See this live on:
vidscrip.com/urojo

Patient well-being is my first priority!

Ureteric Injury

Infrequently ureteric **injuries** can occur with other abdominal surgery ie.

Hysterectomy,
Ovarian mass resection,
Bowel resections,
Sacro Colpo-pxy.


If the injury is below the pelvic brim then a re-implantation with is recommended with Boari flap, otherwise a primary end-to-end anastomosis

Why is it done?

- Strictureing or narrowing of the ureter causing significant hydronephrosis
- Injury to ureter with Urological surgery can cause this: Ureterscopy
- Injury to the ureter from other surgeries: Colorectal, Gynaecological
- This injury if not noted may lead to chronic infection
- End-result is loss of renal function
- An end to end anastomosis can be considered in the mid ureter.
- A Reimplantaion into the bladder with lower ureteric strictures
- A pelvi-ureteric junction repair in higher ureteric strictures
- A trans uretero-ureteric anastomosis joining one ureter to the other where long defects are present
- Renal Auto Transplantation where ureter is completely damaged or
- An ileal ureteric substitution where to whole ureter is damaged

How is it done?

- Laparoscopic or Robotic technique
- Patients will receive a general anaesthesia.
- Prophylactic antibiotics is given.
- The correct ureteric system is identified and marked while you are awake
- This will be mostly a laparoscopic procedure.
- Laparoscopic ports are placed, 1 for camera and 1-2 as working ports.
- The affected ureter is exposed, The defect cut out with a re-anastomosis of a spatulated proximal ureter to a the distal ureter over a ureteric stent.
- Different techniques are used to add more length to the ureter: Boari Flap, Psoas Hitch
- An indwelling catheter is placed.
- A drain is placed.



What next?

- You may be in hospital a few days
- Your catheter will be removed the following day. Or as soon as your urine is clear
- As soon as you are comfortable with no signs of pain and emptying your bladder sufficiently, you will be discharged
- A ward prescription may be issued on your discharge, for your own collection at any pharmacy
- A follow-up appointment will be scheduled for 6 weeks to remove your ureteric stent under local anaesthesia with a Flexible Cystoscopy
- A review with a CT IVP will be scheduled 6 weeks after this to check on the end result of the ureter.
- Any pain or signs of fever require an urgent review
- Don't hesitate to ask Jo if you have any queries
- **DON'T SUFFER IN SILENCE, OR YOU WILL SUFFER ALONE!**

